

Copyright Michael Monagan
You may not distribute my solutions.

Q1 Let S be the sample space for the 5 coin tosses.

So $S = \{ HHHHH, TTTT, \dots \text{etc.} \}$

The outcomes are like binary strings of length 5. So $|S| = 2^5$.

Let A_k be the outcomes with k heads where $0 \leq k \leq 5$

The number of ways to get k heads (and $n-k$ tails) is $\binom{n}{k}$.

$$\text{So } \Pr(X=k) = |A_k|/|S| = \binom{n}{k}/2^n = \binom{5}{k}/32$$

$$\begin{aligned} E[X] &= \sum_{k=0}^5 k \frac{\binom{5}{k}}{32} = \frac{1}{32} [0 \cdot \binom{5}{0} + 1 \cdot \binom{5}{1} + 2 \cdot \binom{5}{2} + 3 \cdot \binom{5}{3} + 4 \cdot \binom{5}{4} + 5 \cdot \binom{5}{5}] \\ &= \frac{1}{32} [0 + 5 + 2 \cdot 10 + 3 \cdot 10 + 4 \cdot 5 + 5 \cdot 1] = 80/32 = 10/4 = 2.5 \end{aligned}$$

Q2 See below.

Q3 Toss $n=5$ balls into $m=5$ bins randomly.

$$\Pr(\text{bin } i \text{ has } 0 \text{ balls}) = \left(\frac{n-1}{n}\right)^m = \left(\frac{4}{5}\right)^5$$

$$\Pr(\text{bin } i \text{ has } 1 \text{ ball}) = n \left(\frac{1}{n}\right) \left(\frac{n-1}{n}\right)^{m-1} = \left(\frac{4}{5}\right)^4$$

$$\Pr(\text{bin } i \text{ has } k \text{ balls}) = \binom{n}{k} \left(\frac{1}{n}\right)^k \left(\frac{n-1}{n}\right)^{m-k} = \binom{5}{k} \left(\frac{1}{5}\right)^k \left(\frac{4}{5}\right)^{5-k}$$

\uparrow # of ways \uparrow k in bin i \uparrow $n-k$ not in bin i

Tabulating the probabilities for $k=0,1,2,3,4,5$ we get

k	0	1	2	3	4	5
$\Pr(X=k)$	$\left(\frac{4}{5}\right)^5$	$\left(\frac{4}{5}\right)^4$	$10 \left(\frac{1}{5}\right)^2 \left(\frac{4}{5}\right)^3$	$10 \left(\frac{1}{5}\right)^3 \left(\frac{4}{5}\right)^2$	$5 \left(\frac{1}{5}\right)^4 \left(\frac{4}{5}\right)$	$\left(\frac{1}{5}\right)^5$

$$\begin{aligned} E[X] &= 1 \cdot \left(\frac{4}{5}\right)^4 + 20 \left(\frac{1}{5}\right)^2 \left(\frac{4}{5}\right)^3 + 30 \left(\frac{1}{5}\right)^3 \left(\frac{4}{5}\right)^2 + 20 \left(\frac{1}{5}\right)^4 \left(\frac{4}{5}\right) + 5 \cdot \left(\frac{1}{5}\right)^5 \\ &= \frac{5 \cdot 4^4 + 20 \cdot 4^3 + 30 \cdot 4^2 + 20 \cdot 4 + 5}{5^5 = 3125} = 1. \quad \checkmark \end{aligned}$$

Q4 $E[X] = n(1 - \frac{1}{n})^m = n\left(\frac{n-1}{n}\right)^m$

$n=10, m=5 \quad E[X] = 5 \cdot 905$

$n=10, m=10 \quad E[X] = 3.487$

$n=10, m=20 \quad E[X] = 1.216$

If we toss 20 balls into 10 bins, on average there will be one empty bin.

If we toss 20 balls into 10 bins, on average there will be one empty bin.

Q5

$$\Pr(\text{bin } i \text{ has one ball}) = m \cdot \left(\frac{1}{n}\right) \left(\frac{n-1}{n}\right)^{m-1} = \frac{m}{n} \left(1 - \frac{1}{n}\right)^{m-1}$$

Let Y be the # of bins with one ball.

Let $Y_i = \begin{cases} 1 & \text{if bin } i \text{ has one ball} \\ 0 & \text{otherwise.} \end{cases}$

$$E[Y] = E[Y_1 + Y_2 + \dots + Y_n] = n E[Y_i]$$

$$= n \cdot (Y_i \cdot \Pr(Y_i))$$

$$= n \cdot 1 \cdot \left(\frac{m}{n}\right) \left(1 - \frac{1}{n}\right)^{m-1} = \boxed{m \cdot \left(1 - \frac{1}{n}\right)^{m-1}}$$

$$= m \left(1 - \frac{1}{n}\right)^n \approx \boxed{m e^{-m/n}} \text{ for large } m.$$

↓ for large n .

For $m=n=10$ $\Pr(Y_i) = \frac{m}{n} \left(1 - \frac{1}{n}\right)^{m-1} = \left(1 - \frac{1}{10}\right)^9 = 0.38742$.

$$E[X] = n \cdot \Pr(Y_i) = m \left(1 - \frac{1}{n}\right)^{m-1} = 10 \left(1 - \frac{1}{10}\right)^9 = 3.8742.$$

$$E[X] \approx m e^{-m/n} = 10 e^{-1} = 3.6789.$$

Q6

$$a_{n+1} = 3a_n - a_{n-1} \text{ with } a_1 = 1, a_2 = 1.$$

$$a_3 = 3a_2 - a_1 = 3 \cdot 1 - 1 = 2$$

$$a_4 = 3a_3 - a_2 = 3 \cdot 2 - 1 = 5$$

$$a_2 = 3a_1 - a_0 \Rightarrow a_0 = 3a_1 - a_2 = 3 \cdot 1 - 1 = 2.$$

Q7.

$$a_n - a_{n-1} = n.$$

$$a_n - a_{n-1} - a_{n-2} = 0$$

Q8.

Let i_n be the # people infected with COVID after n weeks.

We are given $i_n = 3i_{n-1}$ and $i_0 = 2.5$ million.

The general solution is $i_n = A \cdot 3^n$ for $n \geq 1$.

$$i_0 = A = 2.5 \text{ million}$$

So $i_n = 2.5 \cdot 3^n$ million.

$$\text{We need to solve } i_n = 100 = 2.5 \cdot 3^n$$

$$\Rightarrow 3^n = 40 \Rightarrow n = 3.358 \text{ weeks.}$$

Q9.

The # comparisons in line 7 is $n-1$ because the loop is executed from 1 to $n-1$. The # comparisons in line 10 is C_{n-1} because the array to be sorted has length $n-1$. Thus

For $n=1$, $C_1=0$ because of line 4. So

$$C_n = (n-1) + C_{n-1} \quad \text{and} \quad C_1 = 0.$$

$$\Rightarrow C_n = (n-1) + (n-2) + C_{n-2} = \dots = (n-1) + (n-2) + \dots + 1 + C_1$$

$$= (n-1) + (n-2) + \dots + 1 + 0 = \frac{n(n-1)}{2}.$$

This exactly the same # comparisons as Bubblesort.

Q10. Let I, F, and B denote the Italian, French and Brazilian restaurants. For one evening we have $d_1 = 3$ choices. For two evenings we have II, IF, IB, FI, FB, BI, BF, BB so all 6 choices except FF. so $d_2 = 8$.

For three evenings we can't have FF or FF- which rules out 3+2 choices out of $3^3 = 27$ so $d_3 = 22$.

We can eat $\underbrace{I}_1 \underbrace{\leftarrow I \text{ or } B \text{ or } F}_2 \dots \underbrace{\quad}_n$ so d_{n-1} choices

Similarly if we go to B for the first evening. For F

$\underbrace{F}_1 \underbrace{\leftarrow I \text{ or } B}_2 \dots \underbrace{\quad}_n$ so $2d_{n-2}$ choices.

$$\text{Thus } d_n = 2d_{n-1} + 2d_{n-2}$$

$$d_3 = 2d_2 + 2d_1 = 2 \cdot 8 + 2 \cdot 3 = 22$$

$$d_4 = 2 \cdot d_3 + 2 \cdot d_2 = 2 \cdot 22 + 2 \cdot 8 = 44 + 16 = 60$$

$$d_5 = 2 \cdot 60 + 2 \cdot 22 = 164.$$

Q11

$$a_n = \cancel{a_{n-1}} + A_n + B$$

$$\cancel{a_{n-1}} = \cancel{a_{n-2}} + A(n-1) + B$$

$$\cancel{a_{n-2}} = \cancel{a_{n-3}} + A(n-2) + B$$

$$\vdots$$

$$\cancel{a_2} = \cancel{a_1} + A(2) + B$$

$$\cancel{a_1} = 1$$

Adding both sides of these equations we get

$$a_n = 1 + A(n + n-1 + \dots + 2) + (n-1)B = \left[\frac{n(n+1)}{2} - 1 \right] A + (n-1)B + 1.$$

Q12

$$x_n = x_{n-1} + 20x_{n-2} \quad \text{with } x_0 = 5, x_1 = -2.$$

$$\Rightarrow x_n - x_{n-1} - 20x_{n-2} = 0$$

$$\Rightarrow r^2 - r - 20 = (r-5)(r+4) = 0 \Rightarrow r = 5, -4.$$

$$\Rightarrow x_n = A \cdot 5^n + B(-4)^n$$

$$x_0 = A + B = 5$$

$$x_1 = 5A - 4B = -2$$

$$\} \Rightarrow 0 - 9B = -27 \Rightarrow B = 3, A = 2.$$

$$\text{So } x_n = 2 \cdot 5^n + 3(-4)^n.$$

Q2 (b). The Theorem (Linearity of Expectation) says:
If X and Y are random variables and $a \in \mathbb{R}$ Then

(1) $E(aX) = a \cdot E(X)$ and

(2) $E(X+Y) = E(X) + E(Y)$.

So if $Z = X - Y$ then

$$\begin{aligned} E(Z) &= E(X - Y) = E(X + (-Y)) \\ &= E(X) + E(-Y) \text{ by (2)} \\ &= E(X) + (-1)E(Y) \text{ by (1)} \\ &= E(X) - E(Y) = 7 - 3 = 4. \end{aligned}$$

$$Q2 (a) \ E[aX] = \sum_{s \in S} aX(s) \cdot \Pr(s) = a \left(\sum_{s \in S} X(s) \Pr(s) \right) = a \cdot E[X].$$

a is a constant